

#### 4 CBNPC avec réarrangement de ALK

##### 4.1 Traitement par ITK

**-Alectinib** : L'étude ALEX (2) publiée en 2017, randomisant alectinib versus crizotinib en première ligne de traitement montre une nette diminution des progressions cérébrales avec l'alectinib par rapport au crizotinib : 12% versus 45%, (*cause-specific hazard ratio* 0,16,  $p < 0,001$ ), et un taux de réponse intra cérébral de 88% pour l'alectinib vs 50% pour le crizotinib suggérant une bien meilleure efficacité cérébrale de cette molécule. Par ailleurs l'alectinib améliore la survie globale des patients avec des données matures présentées en 2019 à l'ESMO : 34,8 mois dans le bras alectinib versus 10,9 mois dans le bras crizotinib<sup>2</sup>.

**-Brigatinib** : une étude publiée dans le JCO en septembre 2018 a repris 203 patients issus de 2 études de phase 2 (étude ALTA et une autre) avec des MC, le taux de réponse intracérébral des patients ALK traités par brigatinib est entre 46 et 67%, les auteurs concluent à une bonne efficacité du brigatinib au niveau cérébral, qui est meilleure chez les patients traités avec 180mg/j plutôt que 90mg/j (63). L'essai d'enregistrement du Brigatinib en première ligne était à la dose de 90Mg/j pendant 7 jours puis 180mg/j (64).

**L'alectinib et le brigatinib (lorsque son prix sera obtenu dans cette indication) sont donc les ITK à privilégier en première ligne chez les patients avec réarrangement d'ALK avec MC. Leur efficacité au niveau cérébral doit faire privilégier la mise sous ITK par rapport au traitement local qui sera discuté dans un second temps. Le traitement local premier peut quand même être discuté en RCP, notamment en cas de métastase cérébrale unique et/ou symptomatique.**

**-Lorlatinib** : ce TKI de 3ème génération a montré une activité importante au niveau cérébrale chez des patients déjà traités par crizotinib et/ou chimiothérapie et/ou un autre TKI avec des taux de réponse objective intracérébrale de l'ordre de 50 à 60% (65). Actuellement il bénéficie d'une AMM et d'un remboursement chez des patients précédemment traités par alectinib ou ceritinib, ou crizotinib et un autre TKI anti ALK

Dans l'étude CROWN présentée à l'ESMO 2020 (BJ.Solomon et al. LBA2) comparant le lorlatinib en première ligne au crizotinib, 27% des pts présentaient des MC asymptomatiques (65). Le taux de réponse objective intracranienne était de 82% dont 71% de réponse complète pour le lorlatinib contre respectivement 23% et 8% pour le crizotinib (66).

**-Ceritinib** : l'AMM du ceritinib en première ligne permet son utilisation chez les patients ayant des MC au diagnostic mais cette molécule ne doit plus être utilisée en première ligne (cf. référentiel CBNPC). Le taux de réponse cérébrale à 46,3% dans l'étude ASCEND 4 (67), et a donc une efficacité sur le SNC probablement inférieure à l'alectinib (88%). Les résultats préliminaires de l'étude ASCEND 7 montrent un taux de réponse au ceritinib de 51%, taux de contrôle cérébral de 75% chez des patients traités par ceritinib, sans RT ni autre TKI reçu auparavant<sup>3</sup>.

**-Crizotinib** : un traitement par crizotinib en première ligne chez les patients avec réarrangement d'ALK n'est pas recommandé chez les patients porteurs de MC. L'étude PROFILE 1014 (68) montre une amélioration de la PFS et de la survie globale avec le crizotinib en première ligne y compris chez les patients avec des MC.

<sup>2</sup> T.S.K. Mok, A.T. Shaw, R.D. Camidge, S.M. Gadgeel, R. Rosell, R. Dziadziuszko, D. Kim, M. Perol, S. Ou, W. Bordogna, V. Smoljanović, M. Hilton, S. Peters. Final PFS, Updated OS And Safety Data From The Randomised, Phase III ALEX Study Of Alectinib (Alc) Versus Crizotinib (Crz) In Untreated Advanced ALK+ NSCLC. ESMO 2019 – Abs.# 1484PD

<sup>3</sup> L.Q. Chow, F. Barlesi, E.M. Bertino, M.J. van den Bent, H. Wakelee, P.Y. Wen, C. Chiu, S. Orlov, M. Majem, R. Chiari, M. McKeage, C. Yu, F.K. Hurtado, P. Cazorla Arratia, Y. Song, F. Branle, M. Shi, D. Kim. Results Of The Ascend-7 Phase Ii Study Evaluating ALK Inhibitor (Alki) Ceritinib In Patients (Pts) With ALK+ Non-Small Cell Lung Cancer (NSCLC) Metastatic To The Brain. ESMO 2019 – Abs.# 1478O

Cependant il conviendra cependant d'être particulièrement attentif en cas de MC au diagnostic : cette même étude PROFILE 1014 rapporte 43% de progression cérébrale chez les patients avec métastases cérébrales au diagnostic et 22% chez patients sans métastases cérébrales au diagnostic, ce qui suggère un passage de la barrière hémato encéphalique par le crizotinib possiblement insuffisant et inférieur à l'alectinib.

#### 4.2 Progression cérébrale sous ITK

En cas de progression cérébrale sous ITK, la place d'une RTS doit être discutée systématiquement en RCP. En l'absence de données, l'ITK doit être arrêté 48h avant et repris 48 heures après la radiothérapie. Si la progression cérébrale est isolée, asymptomatique et accessible à un traitement local, une poursuite de l'ITK après traitement local cérébral peut être proposée avec surveillance étroite de la maladie cérébrale.

Une radiothérapie des métastases cérébrales accessibles à un traitement en condition stéréotaxique doit systématiquement être discutée, à fortiori quand le nombre de métastase est  $\leq 4$  (score Lung-molGPA 3 à 4, patients avec survies prolongées).

En cas de progression avec MC multiples non accessibles à une RTS, un ITK de deuxième ligne est à privilégier par rapport à une IET du fait de la survie habituellement prolongée de ces patients. L'IET est à éviter ou à discuter au cas par cas en RCP, et la préservation hippocampique doit être réalisée chez ces patients.

**OPTION :** On peut parfois proposer la recherche des mutations de résistance aux ITK d'ALK sur rebiopsie tissulaire ou ADN tumoral circulant. En cas de mise en évidence d'une mutation de résistance, le dossier doit être discuté en RCP de recours, pour inclusion dans un essai si disponible.

#### Recommandations

- Un traitement de première ligne par alectinib ou brigatinib (lorsque le prix sera obtenu dans cette indication) est recommandé.
- En cas de progression cérébrale isolée, une RTS sur les sites cérébraux évolutifs doit être discutée avec la poursuite de l'ITK (qui sera suspendu pendant le temps de la radiothérapie).
- En cas de progression cérébrale et des autres sites métastatiques, le changement d'ITK est recommandé et la recherche des mutations de résistance aux ITK d'ALK sur re-biopsie tissulaire ou ADN tumoral circulant peut être proposée.

## REFERENCES

- Alexander BM, Brown PD, Ahluwalia MS, Aoyama H, Baumert BG, Chang SM, et al. Clinical trial design for local therapies for brain metastases: a guideline by the Response Assessment in Neuro-Oncology Brain Metastases working group. *Lancet Oncol.* 2018;19(1):e33-42.
- Peters S, Camidge DR, Shaw AT, Gadgeel S, Ahn JS, Kim D-W, et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med.* 31 2017;377(9):829-38.
- Chargari C, Dhermain F. Métastases cérébrales d'un cancer bronchique non à petites cellules: du traitement standardisé au traitement personnalisé. *Rev Mal Respir Actual.* 2013;5(5):547-56.
- Sperduto PW, Kased N, Roberge D, Xu Z, Shanley R, Luo X, et al. Summary report on the graded prognostic assessment: an accurate and facile diagnosis-specific tool to estimate survival for patients with brain metastases. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 févr 2012;30(4):419-25.
- Sperduto PW, Yang TJ, Beal K, Pan H, Brown PD, Bangdiwala A, et al. Estimating Survival in Patients With Lung Cancer and Brain Metastases: An Update of the Graded Prognostic Assessment for Lung Cancer Using Molecular Markers (Lung-molGPA). *JAMA Oncol.* 1 juin 2017;3(6):827-31.
- Schellinger PD, Meinck HM, Thron A. Diagnostic accuracy of MRI compared to CCT in patients with brain metastases. *J Neurooncol.* 1999;44(3):275-81.
- Sighvatsson V, Ericson K, Tómasson H. Optimising contrast-enhanced cranial CT for detection of brain metastases. *Acta Radiol Stockh Swed* 1987. nov 1998;39(6):718-22.
- Seute T, Leffers P, ten Velde GPM, Twijnstra A. Detection of brain metastases from small cell lung cancer: consequences of changing imaging techniques (CT versus MRI). *Cancer.* 15 avr 2008;112(8):1827-34.
- Olson JJ, Kalkanis SN, Ryken TC. Congress of Neurological Surgeons Systematic Review and Evidence-Based Guidelines for the Treatment of Adults With Metastatic Brain Tumors: Executive Summary. *Neurosurgery.* 01 2019;84(3):550-2.
- Patchell RA, Tibbs PA, Walsh JW, Dempsey RJ, Maruyama Y, Kryscio RJ, et al. A randomized trial of surgery in the treatment of single metastases to the brain. *N Engl J Med.* 22 févr 1990;322(8):494-500.
- Noordijk EM, Vecht CJ, Haaxma-Reiche H, Padberg GW, Voormolen JH, Hoekstra FH, et al. The choice of treatment of single brain metastasis should be based on extracranial tumor activity and age. *Int J Radiat Oncol Biol Phys.* 1 juill 1994;29(4):711-7.
- Mintz AH, Kestle J, Rathbone MP, Gaspar L, Hugenholtz H, Fisher B, et al. A randomized trial to assess the efficacy of surgery in addition to radiotherapy in patients with a single cerebral metastasis. *Cancer.* 1 oct 1996;78(7):1470-6.
- Latorzeff I, Antoni D, Gaudaire-Josset S, Feuvret L, Tallet-Richard A, Truc G, et al. Radiothérapie des métastases cérébrales. *Cancer/Radiothérapie.* sept 2016;20:S80-7.
- Sahgal A, Aoyama H, Kocher M, Neupane B, Collette S, Tago M, et al. Phase 3 trials of stereotactic radiosurgery with or without whole-brain radiation therapy for 1 to 4 brain metastases: individual patient data meta-analysis. *Int J Radiat Oncol Biol Phys.* 15 mars 2015;91(4):710-7.
- Gerosa M, Nicolato A, Foroni R, Tomazzoli L, Bricolo A. Analysis of long-term outcomes and prognostic factors in patients with non-small cell lung cancer brain metastases treated by gamma knife radiosurgery. *J Neurosurg.* janv 2005;102 Suppl:75-80.
- Motta M, del Vecchio A, Attuati L, Picozzi P, Perna L, Franzin A, et al. Gamma knife radiosurgery for treatment of cerebral metastases from non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys.* 15 nov 2011;81(4):e463-468.
- Fuentes R, Osorio D, Expósito Hernandez J, Simancas-Racines D, Martínez-Zapata MJ, Bonfill Cosp X. Surgery versus stereotactic radiotherapy for people with single or solitary brain metastasis. *Cochrane Database Syst Rev.* 20 2018;8:CD012086.
- Yamamoto M, Serizawa T, Shuto T, Akabane A, Higuchi Y, Kawagishi J, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLKG0901): a multi-institutional prospective observational study. *Lancet Oncol.* avr 2014;15(4):387-95.
- Shuto T, Akabane A, Yamamoto M, Serizawa T, Higuchi Y, Sato Y, et al. Multiinstitutional prospective observational study of stereotactic radiosurgery for patients with multiple brain metastases from non-small cell lung cancer (JLKG0901 study-NSCLC). *J Neurosurg.* 01 2018;129(Suppl1):86-94.
- Le Rhun É, Dhermain F, Noël G, Reyns N, Carpentier A, Mandonnet E, et al. [ANOCEF guidelines for the management of brain metastases]. *Cancer Radiother J Soc Francaise Radiother Oncol.* févr 2015;19(1):66-71.
- Brown PD, Jaeckle K, Ballman KV, Farace E, Cerhan JH, Anderson SK, et al. Effect of Radiosurgery Alone vs Radiosurgery With Whole Brain Radiation Therapy on Cognitive Function in Patients With 1 to 3 Brain Metastases: A Randomized Clinical Trial. *JAMA.* 26 juill 2016;316(4):401-9.
- Mahajan A, Ahmed S, McAleer MF, Weinberg JS, Li J, Brown P, et al. Post-operative stereotactic radiosurgery versus observation for completely resected brain metastases: a single-centre, randomised, controlled, phase 3 trial. *Lancet Oncol.* août 2017;18(8):1040-8.
- Lamba N, Muskens IS, DiRisio AC, Meijer L, Briceno V, Edrees H, et al. Stereotactic radiosurgery versus whole-brain radiotherapy after intracranial metastasis resection: a systematic review and meta-analysis. *Radiat Oncol Lond Engl.* 24 juin 2017;12(1):106.
- Soffiotti R, Kocher M, Abacioglu UM, Villa S, Fauchon F, Baumert BG, et al. A European Organisation for Research and Treatment of Cancer phase III trial of adjuvant whole-brain radiotherapy versus observation in patients with one to three brain metastases from solid tumors after surgical resection or radiosurgery: quality-of-life results. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 janv 2013;31(1):65-72.

25. Brown PD, Ballman KV, Cerhan JH, Anderson SK, Carrero XW, Whitton AC, et al. Postoperative stereotactic radiosurgery compared with whole brain radiotherapy for resected metastatic brain disease (NCCTG N107C/CEC-3): a multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol.* 2017;18(8):1049-60.
26. Ma TM, Grimm J, McIntyre R, Anderson-Keightley H, Kleinberg LR, Hales RK, et al. A prospective evaluation of hippocampal radiation dose volume effects and memory deficits following cranial irradiation. *Radiother Oncol J Eur Soc Ther Radiol Oncol.* 2017;125(2):234-40.
27. Trifiletti DM, Ballman KV, Brown PD, Anderson SK, Carrero XW, Cerhan JH, et al. Optimizing Whole Brain Radiation Therapy Dose and Fractionation: Results From a Prospective Phase 3 Trial (NCCTG N107C [Alliance]/CEC.3). *Int J Radiat Oncol Biol Phys.* 1 févr 2020;106(2):255-60.
28. Rodríguez de Dios N, Couñago F, López JL, Calvo P, Murcia M, Rico M, et al. Treatment Design and Rationale for a Randomized Trial of Prophylactic Cranial Irradiation With or Without Hippocampal Avoidance for SCLC: PREMIER Trial on Behalf of the Oncologic Group for the Study of Lung Cancer/Spanish Radiation Oncology Group-Radiation Oncology Clinical Research Group. *Clin Lung Cancer.* sept 2018;19(5):e693-7.
29. Brown PD, Gondi V, Pugh S, Tome WA, Wefel JS, Armstrong TS, et al. Hippocampal Avoidance During Whole-Brain Radiotherapy Plus Memantine for Patients With Brain Metastases: Phase III Trial NRG Oncology CC001. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 avr 2020;38(10):1019-29.
30. Zindler JD, Bruynzeel AME, Eekers DBP, Hurkmans CW, Swinnen A, Lambin P. Whole brain radiotherapy versus stereotactic radiosurgery for 4-10 brain metastases: a phase III randomised multicentre trial. *BMC Cancer.* 25 juill 2017;17(1):500.
31. Roberge D, Brown PD, Whitton A, O'Callaghan C, Leis A, Greenspoon J, et al. The Future Is Now-Prospective Study of Radiosurgery for More Than 4 Brain Metastases to Start in 2018! *Front Oncol.* 2018;8:380.
32. Robinet G, Thomas P, Breton JL, Léna H, Gouva S, Dabouis G, et al. Results of a phase III study of early versus delayed whole brain radiotherapy with concurrent cisplatin and vinorelbine combination in inoperable brain metastasis of non-small-cell lung cancer: Groupe Français de Pneumo-Cancérologie (GFPC) Protocol 95-1. *Ann Oncol Off J Eur Soc Med Oncol ESMO.* janv 2001;12(1):59-67.
33. Barlesi F, Gervais R, Lena H, Hureaux J, Berard H, Paillot D, et al. Pemetrexed and cisplatin as first-line chemotherapy for advanced non-small-cell lung cancer (NSCLC) with asymptomatic inoperable brain metastases: a multicenter phase II trial (GFPC 07-01). *Ann Oncol Off J Eur Soc Med Oncol ESMO.* nov 2011;22(11):2466-70.
34. Besse B, Lasserre SF, Compton P, Huang J, Augustus S, Rohr U-P. Bevacizumab safety in patients with central nervous system metastases. *Clin Cancer Res Off J Am Assoc Cancer Res.* 1 janv 2010;16(1):269-78.
35. Socinski MA, Langer CJ, Huang JE, Kolb MM, Compton P, Wang L, et al. Safety of bevacizumab in patients with non-small-cell lung cancer and brain metastases. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 nov 2009;27(31):5255-61.
36. Besse B, Le Moulec S, Mazières J, Senellart H, Barlesi F, Chouaid C, et al. Bevacizumab in Patients with Nonsquamous Non-Small Cell Lung Cancer and Asymptomatic, Untreated Brain Metastases (BRAIN): A Nonrandomized, Phase II Study. *Clin Cancer Res Off J Am Assoc Cancer Res.* 15 avr 2015;21(8):1896-903.
37. Brahmer J, Reckamp KL, Baas P, Crinò L, Eberhardt WEE, Poddubska E, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *N Engl J Med.* 9 juill 2015;373(2):123-35.
38. Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med.* 27 sept 2015;
39. Goldman JW, Crino L, Vokes EE, Holgado E, Reckamp K, Pluzanski A, et al. P2.36: Nivolumab (nivo) in Patients (pts) With Advanced (adv) NSCLC and Central Nervous System (CNS) Metastases (mets): Track: Immunotherapy. *J Thorac Oncol Off Publ Int Assoc Study Lung Cancer.* oct 2016;11(10S):S238-9.
40. Hellmann MD, Paz-Ares L, Bernabe Caro R, Zurawski B, Kim S-W, Carcereny Costa E, et al. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. *N Engl J Med.* 28 sept 2019;
41. Goldberg SB, Gettinger SN, Mahajan A, Chiang AC, Herbst RS, Szol M, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol.* juill 2016;17(7):976-83.
42. Gandhi L, Rodríguez-Abreu D, Gadgeel S, Esteban E, Felip E, De Angelis F, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. *N Engl J Med.* 31 mai 2018;378(22):2078-92.
43. Paz-Ares L, Luft A, Vicente D, Tafreshi A, Gümüş M, Mazières J, et al. Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung Cancer. *N Engl J Med.* 22 nov 2018;379(21):2040-51.
44. Rittmeyer A, Barlesi F, Waterkamp D, Park K, Ciardiello F, von Pawel J, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet Lond Engl.* 12 déc 2016;
45. Hendriks LEL, Henon C, Auclin E, Mezquita L, Ferrara R, Audigier-Valette C, et al. Outcome of Patients with Non-Small Cell Lung Cancer and Brain Metastases Treated with Checkpoint Inhibitors. *J Thorac Oncol.* juill 2019;14(7):1244-54.
46. Tallet AV, Dhermain F, Le Rhun E, Noël G, Kirova YM. Combined irradiation and targeted therapy or immune checkpoint blockade in brain metastases: toxicities and efficacy. *Ann Oncol Off J Eur Soc Med Oncol.* 1 déc 2017;28(12):2962-76.
47. Shepard MJ, Xu Z, Donahue J, Eluvathingal Muttikkal TJ, Cordeiro D, Hansen L, et al. Stereotactic radiosurgery with and without checkpoint inhibition for patients with metastatic non-small cell lung cancer to the brain: a matched cohort study. *J Neurosurg.* 26 juill 2019;1-8.

48. Chen L, Douglass J, Kleinberg L, Ye X, Marciscano AE, Forde PM, et al. Concurrent Immune Checkpoint Inhibitors and Stereotactic Radiosurgery for Brain Metastases in Non-Small Cell Lung Cancer, Melanoma, and Renal Cell Carcinoma. *Int J Radiat Oncol Biol Phys.* 15 mars 2018;100(4):916-25.
49. Linskey ME, Andrews DW, Asher AL, Burri SH, Kondziolka D, Robinson PD, et al. The role of stereotactic radiosurgery in the management of patients with newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. *J Neurooncol.* janv 2010;96(1):45-68.
50. Langley RE, Stephens RJ, Nankivell M, Pugh C, Moore B, Navani N, et al. Interim data from the Medical Research Council QUARTZ Trial: does whole brain radiotherapy affect the survival and quality of life of patients with brain metastases from non-small cell lung cancer? *Clin Oncol R Coll Radiol G B.* mars 2013;25(3):e23-30.
51. Mulvenna P, Nankivell M, Barton R, Faivre-Finn C, Wilson P, McColl E, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet Lond Engl.* 22 oct 2016;388(10055):2004-14.
52. Soria J-C, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med.* 18 nov 2017;
53. Ramalingam SS, Vansteenkiste J, Planchard D, Cho BC, Gray JE, Ohe Y, et al. Overall Survival with Osimertinib in Untreated, EGFR-Mutated Advanced NSCLC. *N Engl J Med.* 02 2020;382(1):41-50.
54. Magnuson WJ, Lester-Coll NH, Wu AJ, Yang TJ, Lockney NA, Gerber NK, et al. Management of Brain Metastases in Tyrosine Kinase Inhibitor-Naïve Epidermal Growth Factor Receptor-Mutant Non-Small-Cell Lung Cancer: A Retrospective Multi-Institutional Analysis. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 avr 2017;35(10):1070-7.
55. Miyawaki E, Kenmotsu H, Mori K, Harada H, Mitsuya K, Mamesaya N, et al. Optimal Sequence of Local and EGFR-TKI Therapy for EGFR-Mutant Non-Small Cell Lung Cancer With Brain Metastases Stratified by Number of Brain Metastases. *Int J Radiat Oncol Biol Phys.* 01 2019;104(3):604-13.
56. Lee C-C, Hsu SPC, Lin C-J, Wu H-M, Chen Y-W, Luo Y-H, et al. Epidermal growth factor receptor mutations: association with favorable local tumor control following Gamma Knife radiosurgery in patients with non-small cell lung cancer and brain metastases. *J Neurosurg.* 21 juin 2019;1-8.
57. Reungwetwattana T, Nakagawa K, Cho BC, Cobo M, Cho EK, Bertolini A, et al. CNS Response to Osimertinib Versus Standard Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Patients With Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol Off J Am Soc Clin Oncol.* 28 août 2018;JCO2018783118.
58. Ballard P, Yates JWT, Yang Z, Kim D-W, Yang JC-H, Cantarini M, et al. Preclinical Comparison of Osimertinib with Other EGFR-TKIs in EGFR-Mutant NSCLC Brain Metastases Models, and Early Evidence of Clinical Brain Metastases Activity. *Clin Cancer Res Off J Am Assoc Cancer Res.* 15 oct 2016;22(20):5130-40.
59. Mok TS, Wu Y-L, Ahn M-J, Garassino MC, Kim HR, Ramalingam SS, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med.* 6 déc 2016;
60. Wang X, Xu Y, Tang W, Liu L. Efficacy and Safety of Radiotherapy Plus EGFR-TKIs in NSCLC Patients with Brain Metastases: A Meta-Analysis of Published Data. *Transl Oncol.* oct 2018;11(5):1119-27.
61. Chang EL, Wefel JS, Hess KR, Allen PK, Lang FF, Kornguth DG, et al. Neurocognition in patients with brain metastases treated with radiosurgery or radiosurgery plus whole-brain irradiation: a randomised controlled trial. *Lancet Oncol.* nov 2009;10(11):1037-44.
62. Soffietti R, Kocher M, Abacioglu UM, Villa S, Fauchon F, Baumert BG, et al. A European Organisation for Research and Treatment of Cancer phase III trial of adjuvant whole-brain radiotherapy versus observation in patients with one to three brain metastases from solid tumors after surgical resection or radiosurgery: quality-of-life results. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 janv 2013;31(1):65-72.
63. Camidge DR, Kim D-W, Tiseo M, Langer CJ, Ahn M-J, Shaw AT, et al. Exploratory Analysis of Brigatinib Activity in Patients With Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer and Brain Metastases in Two Clinical Trials. *J Clin Oncol Off J Am Soc Clin Oncol.* 10 2018;36(26):2693-701.
64. Camidge DR, Kim HR, Ahn M-J, Yang JC-H, Han J-Y, Lee J-S, et al. Brigatinib versus Crizotinib in ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med.* 22 nov 2018;379(21):2027-39.
65. Solomon BJ, Besse B, Bauer TM, Felip E, Soo RA, Camidge DR, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study. *Lancet Oncol.* déc 2018;19(12):1654-67.
66. Shaw AT, Bauer TM, de Marinis F, Felip E, Goto Y, Liu G, et al. First-Line Lorlatinib or Crizotinib in Advanced ALK-Positive Lung Cancer. *N Engl J Med.* 19 nov 2020;383(21):2018-29.
67. Soria J-C, Tan DSW, Chiari R, Wu Y-L, Paz-Ares L, Wolf J, et al. First-line ceritinib versus platinum-based chemotherapy in advanced ALK-rearranged non-small-cell lung cancer (ASCEND-4): a randomised, open-label, phase 3 study. *Lancet Lond Engl.* 4 mars 2017;389(10072):917-29.
68. Solomon BJ, Mok T, Kim D-W, Wu Y-L, Nakagawa K, Mekhail T, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med.* 4 déc 2014;371(23):2167-77.
69. Pereira I, Slotman B, Rusthoven CG, Katz MS, Simcock R, Saeed H. Stereotactic radiosurgery (SRS) - A new normal for small cell lung cancer? *Clin Transl Radiat Oncol.* nov 2020;25:10-5.
70. Horn L, Mansfield AS, Szczesna A, Havel L, Krzakowski M, Hochmair MJ, et al. First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer. *N Engl J Med.* 25 sept 2018;

71. Paz-Ares L, Dvorkin M, Chen Y, Reinmuth N, Hotta K, Trukhin D, et al. Durvalumab plus platinum–etoposide versus platinum–etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial. *The Lancet*. oct 2019;S0140673619322226.
72. Horn L, Mansfield AS, Szczesna A, Havel L, Krzakowski M, Hochmair MJ, et al. First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer. *N Engl J Med*. 6 déc 2018;379(23):2220-9.